

Transcript Details

This is a transcript of a continuing medical education (CME) activity. Additional media formats for the activity and full activity details (including sponsor and supporter, disclosures, and instructions for claiming credit) are available by visiting:

<https://reachmd.com/programs/cme/connective-tissue-diseases-ctd-identifying-patients-with-ph/14400/>

Released: 07/29/2022

Valid until: 07/29/2023

Time needed to complete: 1h 16m

ReachMD

www.reachmd.com

info@reachmd.com

(866) 423-7849

Connective Tissue Diseases (CTD): Identifying Patients With PH

Announcer:

Welcome to CME on ReachMD. This episode is part of our MinuteCME curriculum.

Prior to beginning the activity, please be sure to review the faculty and commercial support disclosure statements as well as the learning objectives.

Dr. Krasuski:

Connective Tissue Diseases. Identifying Patients with Pulmonary Hypertension. Patients with systemic sclerosis can have different reasons for their pulmonary hypertension. As you can see, there's tremendous crossover here. They can have Group 1, or PAH. There can be patients with significant interstitial lung disease, and those folks fall predominantly into Group 3. They can also develop Group 2 disease because it's well known that myocardial problems also can occur. So there's a tremendous overlap that eventually leads to right heart failure.

Here is an example of the different types of pulmonary hypertension in systemic sclerosis. On the left you can see a patient with interstitial lung disease. In this case, it's much more diffuse than limited. Their very abnormal lung architecture is present. There's reduced vital capacity and these patients are chronically hypoxic. There's also normal to moderate PA pressure elevation. So these are patients who predominantly have an ILD component with some pulmonary hypertension. On the right, on the other hand, is somebody with pulmonary arterial disease. In this case limited considerably is greater than the diffuse type of disease. There is normal or minimally abnormal lung architecture. There's mild basilar fibrosis often. There's very low DLco and a normal vital capacity. And there's evidence of right ventricular dysfunction. The heart typically becomes significantly greater in size and there's elevated PA pressures. In this case, you have rather significant pulmonary hypertension.

Announcer:

You have been listening to CME on ReachMD. This activity is jointly provided by Global Learning Collaborative (GLC) and TotalCME, Inc. and is part of our MinuteCME curriculum.

To receive your free CME credit, or to download this activity, go to ReachMD.com/CME. Thank you for listening.