

### Transcript Details

This is a transcript of an educational program accessible on the ReachMD network. Details about the program and additional media formats for the program are accessible by visiting: <https://reachmd.com/programs/medical-industry-feature/changing-our-view-of-acute-myeloid-leukemia-in-older-adults/11612/>

### ReachMD

www.reachmd.com  
info@reachmd.com  
(866) 423-7849

---

### Changing Our View of Acute Myeloid Leukemia in Older Adults

#### ReachMD Announcer:

Welcome to ReachMD. The following program, "Changing Our View of Acute Myeloid Leukemia in Older Adults" is developed and sponsored by AbbVie. This activity is intended for United States and Puerto Rico health care professionals only.

The US Medical Affairs department of AbbVie Inc. is the sole author and copyright owner of this presentation and has paid ReachMD to host this presentation. AbbVie is solely responsible for all written and oral content within this presentation. ©2020 AbbVie Inc. All rights reserved.

The following speaker has received compensation from the US Medical Affairs department of AbbVie Inc. to prepare and present the following information and is speaking on behalf of AbbVie.

#### Dr. Erba:

AML is the most common acute leukemia that affects adults, representing 1.2 percent of all new cancer cases in the United States. Survival among younger patients with AML has improved over the past decade; however, older patients continue to have a poor prognosis.

Hello! I am Dr. Harry Erba, director of the Leukemia Program and Medical Director of the Hematologic Malignancies Inpatient Service at Duke University in Durham, North Carolina. And, in this episode, I will be discussing the changing standards of care in acute myeloid leukemia, or AML.

Studies from real-world practice and experimental clinical trials have demonstrated that age and frailty are the primary predictors of outcome among patients with AML. Therefore, proper assessment of a patient's fitness has represented a critical step in the therapeutic decision-making process in AML.

In the past, we focused on curing patients who are eligible for intensive chemotherapy. We have tried to achieve complete responses without measurable residual disease, that is, MRD negativity, in these patients because we know they have a higher chance of cure without relapse. In fact, those patients considered to be eligible for intensive induction chemotherapy have high rates of complete remission, whereas patients considered to be ineligible had much lower rates of complete remission with these choices of less intensive chemotherapy.

The majority of patients with AML do not receive intensive chemotherapy; therefore, we must change the way we think about older patients with AML. Because only 15 to 20% of older patients were expected to achieve complete remission with less intensive chemotherapy, and less intensive chemotherapy had not improved overall survival compared with best supportive care alone, many did not receive any treatment other than best supportive care and/or hospice care.

But today, as a result of an increased understanding of AML pathogenesis, agents exist that target the disease on a molecular level, and do not have the same toxicities as intensive cytotoxic therapy, offering alternative therapeutic interventions for older patients.

#### ReachMD Announcer:

This program was brought to you by AbbVie. If you missed any part of this discussion or to find others in this series, visit [ReachMD.com/SpotlightOn](https://ReachMD.com/SpotlightOn). This is ReachMD. Be part of the knowledge.

BCL2-US-00036-MC, Version 1.0  
Approved: November 2020

